

Digestive Disease Associates of Rockland, PC
974 Route 45, Pomona NY 10970
845-354-3700

FLEXIBLE SIGMOIDOSCOPY PREPARATION

You have been scheduled for a flexible sigmoidoscopy with Digestive Disease Associates of Rockland. This is an examination of the lower portion of your large intestine (colon). Under sedation, a long flexible tube called a colonoscope will be inserted into your rectum and passed through your colon. Your colon will be examined in detail. Additional procedures may be performed such as taking tissue samples (biopsies) or removing polyps.

Please read all the instructions at least ONE WEEK before your scheduled procedure date so you can be adequately prepared for this procedure. A thorough cleansing of the colon is essential and the examination is most successful if you follow the directions for preparation completely. Improper cleansing may result in rescheduling your procedure. If you have any questions, please call our office at 845-354-3700. Additional information can also be found on our website at www.myddar.com.

To ensure you are comfortable and relaxed during your procedure, intravenous sedation may be given. **YOU MAY NOT DRIVE HOME AFTER YOU RECEIVE SEDATION.** You must have a responsible adult escort with a valid driver's license who will be able to drive you home after your procedure. **You may not go home alone** in a taxi, shuttle van or bus, as the drivers will not be responsible for you. You may not drive until the day after your procedure. We advise you to take the entire procedure day off from work.

Our highest priority is performing the most careful and safest possible examination for each and every patient. Although we strive to remain on schedule, please understand that performing certain procedures may take longer in some patients than in others. As a result, your procedure may not be performed at the exact time you were scheduled. We ask for your patience and advise that you allow at least **4 hours** for your visit.

SPECIAL CONSIDERATIONS

Please **bring a list of your current medications**, along with dosages, with you the day of your procedure.

- You should continue to take most of your medications, including your **BLOOD PRESSURE** and **HEART MEDICATIONS**, without interruption, including on the morning of your procedure.
- **Blood thinning medications** (such as Coumadin, warfarin, Plavix, Pradaxa, Xarelto, Effient, Lovenox, Aggrenox) **MAY** need to be discontinued for **1 to 10 days** prior to your procedure, depending on the circumstances. **Please inform us** if you are taking these medications. We recommend that you speak with your cardiologist or primary physician regarding stopping your blood thinners.
- Stop **ASPIRIN, ADVIL, MOTRIN, ALEVE, IBUPROFEN** or any other anti-inflammatory drugs 7 days before your procedure. If needed, you may take **TYLENOL** only.
- **Iron pills, Vitamin E, Gingko biloba:** Stop taking 7 days prior to your procedure.
- If you are **DIABETIC**, check with your primary physician regarding taking your insulin or oral diabetic medications. In general, you should not take any oral diabetic medications the day before or the day of your procedure. Speak with your primary physician if you are on insulin regarding how to adjust the insulin dosage.
- **All women under 50 years** of age must have a urine pregnancy test performed in our office, prior to your procedure. Please be prepared to give a urine sample when you are called in to the procedure area.

SIGMOIDOSCOPY PREP INSTRUCTIONS

PURCHASE THESE ITEMS AHEAD OF TIME (no prescription is required):

- Dulcolax laxative tablets (2 tablets)
- Two (2) Fleets Rectal Enemas

THE DAY BEFORE YOUR SIGMOIDOSCOPY:

- Take 2 Dulcolax tablets at 7 PM.
- You may have regular food all day, including dinner.

THE DAY OF YOUR SIGMOIDOSCOPY:

- **You may not have anything to eat or drink the entire day**, until after your procedure is completed.
- If you take **prescription medications**, you may take them in the morning with a small sip of water. Continue all your usual prescription medications - please be sure to take any blood pressure or heart medications the morning of the test with a small sip of water.

TWO HOURS BEFORE YOUR PROCEDURE TIME:

- Remove the cap from the Fleets enema bottle
- Bend over; you do not need to lie down
- Insert the lubricated tip of the enema into your rectum.
- Squeezed the entire contents of the enema bottle into your rectum.
- Sit on the toilet; hold the enema as long as you can, or up to 5 min. before releasing.
- Repeat the above steps with the second Fleets enema.